

# Exact results for a one dimensional $t - J$ model with impurity

P.-A. Bares

*Institut Laue-Langevin, B.P. 156x, F-38042 Grenoble Cedex, France*

(February 1, 2008)

## Abstract

We propose a simple idea to construct 1D integrable models with impurities. We illustrate the strategy for a supersymmetric  $t - J$  Hamiltonian in considerable detail. The impurity comprises the local deformation of the hopping and exchange integrals as well as a three-body charge-current interactions on neighboring sites. We explore the thermodynamic properties of the system at low and high temperatures, and obtain results that are beyond boundary conformal-field theory.

PACS1993: 71.27.+a, 71.30.+h, 05.30.Fk

Recently there has been new developments in the theory of impurities in Fermi and Luttinger liquids. This renewal of interest in the quantum impurity problems has been stimulated in part by the search for non-Fermi-liquid fixed-points beyond one space dimension and its plausible relevance to the theory of heavy fermions and high temperature superconductors. In this context, Affleck, Ludwig and collaborators have used the (multi-channel) Kondo problem as a laboratory to test, develop and extend significantly the techniques of conformal field theory in the presence of boundaries [1,2]. These powerful methods allow to obtain single-particle as well as thermodynamic properties near the critical point. Using simpler bosonization and renormalization group techniques, Kane and Fisher have shown that a potential scatterer embedded in a Luttinger liquid is driven to a strong-coupling fixed point by the electron-electron repulsive interactions [3]. Emery and Kivelson were able to map the two channel Kondo problem to a resonant-level Hamiltonian which could be solved exactly at a particular point [4]. Fabrizio and Gogolin have generalized the Emery-Kivelson solution to the isotropic four channel and together with Nozières to the anisotropic multi-channel Kondo problem [5,6]. Combining bosonization and conformal field theory results, Sengupta and Georges have proposed a solution for a two-channel two-impurity Kondo Hamiltonian [7]. Significant progress in a different direction has also been achieved by Ghoshal and Zamolodchikov. These authors were able to solve exactly integrable field theories with boundaries [8]. Interesting applications of these results have been performed by Fendley et al. and Tsvelick [9].

Despite this important progress, the problem of an (few) impurity(ies) (potential, magnetic or boths) embedded in a strongly correlated system is not well understood. There is a notable exception: a spin  $S > 1/2$  impurity in a spin  $1/2$  Heisenberg chain solved many years ago by Andrei and Johannesson [10] and generalized to arbitrary spins by Lee and Schlottmann, and Schlottmann [11]. Very recently, it was argued in reference [12] that these models correspond to non-generic fixed points. In the present letter, we want to explore a new idea that allows us to construct exactly solvable models with a (few) defect(s). Short range interactions between the impurities can also be incorporated. There is a prize to pay,

however, namely some terms in the Hamiltonian are difficult to justify on physical grounds. Their presence is required by formal reasons (integrability) and should be generically irrelevant to the low-energy properties of the system. The idea is simple and can be explained as follows. Consider an integrable model that can be formulated within the framework of the Quantum Inverse Scattering Method (QISM) [14]. In the QISM, the Hamiltonian of the model is written as the logarithmic derivative of an homogeneous transfer matrix at a special value of the spectral parameter (an infinite number of conserved kinematic quantities can be generated by performing successive derivatives of the logarithm of the transfer matrix). In general, however, the transfer matrix may depend on a set of (unequal) parameters. In the statistical-mechanics language this corresponds to an inhomogeneous vertex model. Define the Hamiltonian as the logarithmic derivative of the corresponding inhomogeneous transfer matrix. Provided we restrict to an (few) inhomogeneity(ies), we have a model with a (few) defect(s) that is exactly integrable by construction. We have performed preliminary calculations in a variety of models: spinless fermions, spin chains and strongly correlated systems. A general discussion will appear elsewhere [15].

In this work, we want to illustrate the idea by focusing on one particular model that is a prototype for strongly correlated electrons, i.e., the  $t - J$  model. In the absence of impurity, the  $t - J$  model is known to be solvable at the supersymmetric point (away from the integrable point, much is also known through numerical studies, see ref. [16]) and various properties of the model have been investigated in the recent past [17–19]. To construct a  $t - J$  model with impurity, we follow the strategy proposed above and arrive at the Hamiltonian:  $H = H_0 + H_I$ , where  $H_0 = \sum_{n=1}^L H_{nn-1}$  denotes the usual  $t - J$  Hamiltonian,

$$H_{nn+1} = \mathcal{P} \left\{ - \sum_{\sigma} (c_{n\sigma}^{\dagger} c_{n+1\sigma} + h.c.) + J (\mathbf{S}_{n+1} \cdot \mathbf{S}_n - \frac{\hat{n}_n \hat{n}_{n+1}}{4}) + \hat{n}_n + \hat{n}_{n+1} \right\} \mathcal{P} , \quad (1)$$

where  $\mathcal{P}$  projects out the doubly occupied sites,  $\hat{n}_n = \sum_{\sigma} c_{n\sigma}^{\dagger} c_{n\sigma}$ ,  $J = 2$  and periodic boundary conditions have been assumed. The impurity contribution can be written as

$$H_I = g(H_{m-1m+1} - H_{mm-1} - H_{mm+1}) + i\sqrt{g(1-g)}[H_{m-1m+1}, H_{mm-1}] , \quad (2)$$

where the coupling constant  $0 \leq g \leq 1$  (this restriction follows from the hermiticity of  $H$ ) and  $m$  denotes the center of the defect that involves, besides  $m$ , its two adjacent sites  $m \pm 1$ . The first term on the r.h.s. of (2) enhances antiferromagnetic correlations between  $m \pm 1$  sites while ferromagnetic alignment is favored between  $m$  and its two adjacent sites. The second term corresponds to a three-body interaction,

$$\begin{aligned} \mathcal{P} \{ & (1 - n_m) \sum_{\sigma} (j_{m+1m-1}^{\sigma\sigma} + h.c.) - (1 - n_{m-1}) \sum_{\sigma} (j_{m+1m}^{\sigma\sigma} + h.c.) - (1 - n_{m+1}) \sum_{\sigma} (j_{mm-1}^{\sigma\sigma} + h.c.) \\ & + \sum_{\sigma\sigma'} (c_{m\sigma}^{\dagger} c_{m\sigma'} j_{m+1m-1}^{\sigma'\sigma} + h.c.) - \sum_{\sigma\sigma'} (c_{m-1\sigma}^{\dagger} c_{m-1\sigma'} j_{m+1m}^{\sigma'\sigma} + h.c.) - \sum_{\sigma\sigma'} (c_{m+1\sigma}^{\dagger} c_{m+1\sigma'} j_{mm-1}^{\sigma'\sigma} + h.c.) \\ & - \sum_{\sigma\sigma'} (c_{m\sigma}^{\dagger} c_{m\sigma'} J_{m-1m+1}^{\sigma\sigma'} + h.c.) \} \mathcal{P} \quad , \end{aligned} \quad (3)$$

with  $j_{ab}^{\sigma\sigma'} = i c_{a\sigma}^{\dagger} c_{b\sigma'}$  and  $J_{ab}^{\sigma\sigma'} = i \sum_{\lambda} (c_{a\sigma'}^{\dagger} c_{a\lambda}) (c_{b\lambda}^{\dagger} c_{b\sigma})$ . In (3), we have three types of terms: (i) a local charge current flowing between the sites  $m$ ,  $m \pm 1$  provided one of them is empty; (ii) assisted spin-flip currents; (iii) pure assisted spin-currents. At  $g = 1$ , the three body term is absent and the model behaves as a chain of  $L - 1$  sites with no impurity present.

The eigenvalue problem associated to  $H$  is solved by the graded version of the QISM (see ref. [20]). For convenience, we choose the Sutherland representation (see discussion in [18]). The formal solution is casted into the form of the so-called Bethe ansatz equations,

$$L p_{\alpha}(\lambda_j^{\alpha}) + \delta_{\alpha}(\lambda_j^{\alpha}) = 2\pi I_j^{\alpha} - \sum_{\beta=c,s} \sum_{k=1}^{N_{\beta}} \phi_{\alpha\beta}^0(\lambda_j^{\alpha} - \lambda_k^{\beta}) \quad , \quad (4)$$

where  $p_{\alpha}(\lambda_j^{\alpha})$ , for  $\alpha = c, s$ , denotes the bare momentum with  $p_s(x) = 2 \tan^{-1}(2x)$  and  $p_c(x) = 0$ ,  $\phi_{\alpha\beta}^0(x)$  is the bare phase shift for the scattering of the  $\alpha$  with the  $\beta$  pseudoparticles ( $\phi_{ss}^0(x) = -2 \tan^{-1}(x)$ ,  $\phi_{sc}^0(x) = \phi_{cs}^0(x) = 2 \tan^{-1}(2x)$ ,  $\phi_{cc}^0(x) = 0$ ) while  $\delta_{\alpha}(x)$  denotes the bare phase shift for the scattering of the  $\alpha$  pseudoparticle with the impurity, i.e.,  $\delta_s(x) = 2[\tan^{-1}(2[x-\theta]) - \tan^{-1}(2x)]$  and  $\delta_c(x) = 0$ . The parameter  $\theta$  is defined by  $\theta = (g/(1-g))^{1/2}$ . As usual the quantum numbers  $I_j^{\alpha}$  are integers or half-odd integers depending on the number of sites  $L$ , the number of particles  $N$  and the magnetization  $M$  in the system. The ground-state (of finite momentum) and part of the low-energy spectrum are parametrized by real roots in equation (4) (see ref. [18]).

Since we are interested in thermodynamic properties, we shall directly work at finite temperature with the thermodynamic potential per unit length  $\Omega(\mu, H, T)$ , where  $\mu$  denotes the chemical potential,  $H$  the external magnetic field and  $T$  the temperature. At finite  $T$ , we have to consider all possible solutions of the set of equations (4). It turns out that the rapidities associated to the s-pseudoparticles  $\lambda_j^s$  cluster into strings in much the same way as they do in the Heisenberg chain [21] while the “charge” rapidities  $\lambda_j^c$  remain real. Following Yang and Yang [22], we write down the thermodynamic potential and minimize with respect to the distribution of roots of the set of equations (4). After some algebra (see for example [23]), the thermodynamic potential per unit length, i.e., the pressure, is found, up to an irrelevant constant,

$$\Omega = -\mu - h + \frac{1}{\beta} \sum_{n=1}^{\infty} \int dx \left[ a_n(x) + \frac{1}{L} (a_n(x - \theta) - a_n(x)) \right] \log [1 - f(\epsilon_{n,s}(x))] , \quad (5)$$

where  $h = \mu_B H$ ,  $\mu_B$  is the Bohr magneton,  $\beta = (k_B T)^{-1}$ ,  $a_n(x) = (2\pi)^{-1} dp_n(x)/dx$  with  $p_n(x) = 2 \tan^{-1}(2x/n)$ .  $\Omega$  naturally separates into a bulk contribution of  $O(1)$  and an impurity contribution of  $O(1/L)$ . The function  $f(\epsilon_{sn}) = (1 + e^{\beta \epsilon_{sn}})^{-1}$  coincides with the Fermi distribution function in the low-temperature limit while it differs from the former at finite  $T$  where the quasi-energies  $\epsilon_{ns}(x)$  are  $T$ -dependent. The latter are determined from the integral equations

$$\begin{aligned} \beta \epsilon_{sn}(x) &= s \star \log (f(\epsilon_{sn-1}) f(\epsilon_{sn+1})) (x) \\ &+ \delta_{n,1} [s \star \log (1 - f(\epsilon_c)) (x) - 2\pi \beta J s(x)] , \end{aligned} \quad (6a)$$

$$\beta \epsilon_c(x) = \beta \mu - \beta h + \sum_n a_n \star \log (1 - f(\epsilon_{sn})) (x) , \quad (6b)$$

where  $s(x) = 1/2 \cosh(\pi x)$ ,  $\star$  denotes a convolution and the boundary conditions,  $\lim_{n \rightarrow \infty} (\epsilon_{ns}/n) = h$ , have been imposed. Notice that (6a) and (6b) do not depend on the impurity because the contribution of the latter to the free energy vanishes in the variation with respect to the densities of roots.

In general, the set of coupled equations (6a) and (6b) are difficult to solve analytically. In the following, we solve these equations in the low- and high- $T$  limits. We use techniques

similar to those developped by Takahashi [24]. We first discuss the high temperature region ( $\beta J \gg 1$ ), i.e., the weak coupling limit. To lowest order in  $\beta J$ , the pseudo-Fermi distribution functions  $f(\epsilon_c)$  and  $f(\epsilon_{sn})$  are constants, and (6a) and (6b) simplify to  $f(\epsilon_{sn}) = F_n^{-2}$  and  $f(\epsilon_c) = 1 - F_0^2$ , where  $F_n = (bz^n - b^{-1}z^{-n})/(b - b^{-1})$  with  $z = e^{-\beta h}$  and  $b = [(1 + ze^{\beta\mu})/(1 + z^{-1}e^{\beta\mu})]^{1/2}$ . Substitution of this result into the thermodynamic potential (5) leads to:  $\Omega_0 = -\beta^{-1} \log(1 + 2e^{\beta\mu} \cosh(\beta h))$ . We note that the impurity does not contribute in this limit.  $\Omega_0$  is just the thermodynamic potential per unit length for a lattice where each site has three possible states [17]. In this case the entropy, specific heat, etc. are easy to interpret. To higher order in  $\beta$ , the quasi-energies are no longer constants and we expand them as  $\beta\epsilon_{sj}(x) = \beta\epsilon_{sj}^0(x) + \sum_{n=1}^{\infty} (\pi\beta J)^n \zeta_{sj}^{(n)}(x)$ , and  $\beta\epsilon_c(x) = \beta\epsilon_c^0(x) + \sum_{n=1}^{\infty} (\pi\beta J)^n \zeta_c^{(n)}(x)$ , where  $\beta\epsilon_{sn}^0 = \log(F_{n-1}F_{n+1})$  and  $\beta\epsilon_c^0 = \log(F_0^2/(1 - F_0^2))$ . Below we discuss the lightly doped Mott-Hubbard insulator. Near half-filling, we have solved for  $\zeta_{sj}^{(1)}(x)$  and  $\zeta_c^{(1)}(x)$  and their Fourier transforms are  $\hat{\zeta}_{sn}^{(1)}(\omega) = -(F_0/F_1) \left[ (F_n/F_{n-1})e^{-n|\frac{\omega}{2}|} - (F_n/F_{n+1})e^{-(n+2)|\frac{\omega}{2}|} \right]$ , and  $\hat{\zeta}_c^{(1)}(\omega) = -(1/F_1^2)e^{-|\omega|}$ . Substituting this into (5), leads to the thermodynamic potential  $\Omega = \Omega_0 + \Omega_1 + O((\beta J)^2)$ , where  $\Omega_1 = \Omega_1^h + \Omega_1^i/L = -(J/F_1^2)(1 - g/L)$ . From the latter we infer immediately that

$$\frac{\Delta S^i}{\Delta S^h} = \frac{\Delta C_v^i}{\Delta C_v^h} = \frac{\Delta M^i}{\Delta M^h} = \frac{\Delta \chi_s^i}{\Delta \chi_s^h} = \frac{\Delta \chi_c^i}{\Delta \chi_c^h} = -g, \quad (7)$$

where  $\Delta S^{i,h}$ ,  $\Delta C_v^{i,h}$ ,  $\Delta M^{i,h}$ ,  $\Delta \chi_s^{i,h}$ ,  $\Delta \chi_c^{i,h}$  denote the  $O(\beta J)$  corrections to the impurity (host) entropy, specific heat, magnetization, spin susceptibility, and compressibility, respectively. The simplicity of this formula is remarkable. It expresses the fact that in the weak coupling limit, the impurity “mimics” the host chain. Notice that at  $g = 1$ , the expression for the thermodynamic potential density (remember to multiply by  $L/(L - 1)$ ) reduces to that of a chain of  $L - 1$  sites. We have evaluated explicitly the entropy, specific heat, etc. to  $O(\beta J)$  as a function of  $\mu$ ,  $h$  and  $T$ . For brevity, we mention below only the simplest results [15]. Inverting the equation that defines the density  $\rho = 1 - \delta$ , ( $\delta$  = hole density) as function of  $\mu$ , yields for  $\delta \ll 1$

$$\mu = -\beta^{-1} \left[ \log \delta + \log(1 + \delta) + \log(2 \cosh(\beta h)) + \beta J \left( 1 - \frac{g}{L} \right) \right] + O(\delta^3, (\beta J)^2). \quad (8)$$

In the Sutherland representation, the Heisenberg chain is obtained for  $\mu \rightarrow \infty$  (remember that  $\mu$  plays the role of the chemical potential for the holes). Substituting (8) back into the entropy per unit length, at  $h = 0$ , we have

$$\begin{aligned} \frac{S}{k_B} &\approx \log 2 + \delta(1 - [\log \delta + \log 2]) - \frac{\delta^2}{2} \\ &+ \frac{\beta J}{2} \left(1 - \frac{g}{L}\right) \{\delta[\log \delta + \log 2] + \delta^2(1 - 2[\log \delta + \log 2])\} + O(\delta^3, (\beta J)^2) , \end{aligned} \quad (9)$$

The various contributions to the entropy are easy to interpret: (i) the entropy of free spins 1/2 (of density  $1 - \delta$ ); (ii) the entropy of free holes; (iii) a reduction of the entropy due to the exchange correlations ( $J > 0$  and  $\log \delta < -\log 2$  for  $\delta \ll 1$ ); (iv) an increase in entropy due to the impurity that depletes locally the exchange. The specific heat in zero field near half-filling behaves as

$$C_v = k_B \delta [\log \delta + \log 2] \left[ [\log \delta + \log 2] - \beta J \left(1 - \frac{g}{L}\right) \left(1 + \frac{1}{2} [\log \delta + \log 2]\right) \right] + O(\delta^3, (\beta J)^2) , \quad (10)$$

The magnetization is finite in an external field and near half-filling takes the form

$$\frac{M}{\mu_B} = (1 - \delta) \tanh(\beta h) - \beta J \left(1 - \frac{g}{L}\right) (1 - \tanh^2(\beta h))(1 - 3\delta + 3\delta^2) + O(\delta^3, (\beta J)^2) , \quad (11)$$

i.e., the magnetization is suppressed by (i) the doping that reduces the number of spins; (ii) the antiferromagnetic correlations due to the exchange. To next order, the defect enhances the magnetization locally, while the doping renormalizes down the effective exchange between nearest neighbors. The  $\delta^2$  term can be interpreted as an attractive interaction between the holes that effectively increases the renormalized exchange and thereby penalizes the magnetic state. We have also evaluated the charge and spin susceptibilities to  $O(\delta^2)$ . The general expressions are rather complex and will be discussed elsewhere [15].

We now pass to the low temperature region, i.e., strong coupling regime. We note that the quasi-energies  $\epsilon_{sn}(x) > 0$  for  $n \geq 2$ ,  $h > 0$  (compare with ref. [24]) and so as  $T \rightarrow 0$ , only  $\epsilon_{s1}(x) = \epsilon_s(x)$  and  $\epsilon_c(x)$  survive in equations (6a) and (6b) [25]. At  $T = 0$ , the integral equations determining  $\epsilon_s(x)$  and  $\epsilon_c(x)$  coincide with those of ref. [18]. We have explicitly

evaluated the distribution of roots as well as the quasi-energies in zero field near half-filling [15]. The first non-trivial corrections to the spinon and holon quasi-energies appear to third order in the density of holes  $\delta_0$  ( $\delta = \delta_0 + \delta_i/L$ ), i.e.,

$$\epsilon_s(x) \approx -\frac{\pi J}{2 \cosh(\pi x)} \left[ 1 + \delta_0^3 \left( \frac{\pi}{2 \log 2} \right)^3 \left( \frac{\zeta(3)}{\pi} - \frac{5\pi \log 2}{16} [1 - 2 \tanh^2(\pi x)] \right) \right] , \quad (12a)$$

$$P_s(x) \approx \frac{\pi}{2} - \tanh^{-1}(\sinh(\pi x))(1 + \delta_0) + \frac{\delta_0^3}{3} \left( \frac{\pi}{2 \log 2} \right)^3 \left[ \pi \log 2 \frac{\tanh(\pi x)}{\cosh(\pi x)} - \frac{3\zeta(3)}{2\pi} + \left( \frac{2 \log 2}{\pi} \right)^3 \right] , \quad (12b)$$

$$\epsilon_c(x) \approx \pi J \left[ \frac{\log 2}{\pi} - R(x) + \delta_0^2 \frac{3\zeta(3)}{4\pi} + \delta_0^3 \left( \frac{3\zeta(3)}{2\pi} \left( \frac{\pi}{2 \log 2} \right)^2 - \frac{\zeta(3)}{\pi} R(x) - \frac{5 \log 2}{16\pi} R''(x) \right) \right] , \quad (13a)$$

$$P_c(x) \approx 2\pi F(x) \left[ 1 + \delta_0 - \delta_0^3 \frac{\zeta(3)}{2\pi} \left( \frac{\pi}{2 \log 2} \right)^3 \right] - \delta_0^3 \frac{\pi}{3} \left( \frac{\pi}{2 \log 2} \right)^2 R'(x) , \quad (13b)$$

where  $P_{c(s)}(x)$  denotes the momentum of the  $c(s)$ -type excitation relative to the ground-state,  $\zeta(3)$  is the Riemann zeta function,  $R(x) = \Re [\Psi(1 + ix/2) - \Psi([1 + ix]/2)] / 2\pi$  with the definition  $\Psi(x) = d \log \Gamma(x) / dx$ .  $R'(x)$  and  $R''(x)$  denotes the first and second derivatives of  $R(x)$  while  $2\pi F(x) = (\log \pi)/2 + \log (\Gamma(1 + ix/2)/\Gamma([1 + ix]/2))$ . The  $O(\delta_0^2)$  term in  $\epsilon_c(x)$  is due to a chemical potential correction, i.e.,  $\mu = J[\log 2 - 3\zeta(3)\delta_0^2/4] + O(\delta^4)$ , that shifts the bottom of the holon spectrum upwards. From (12a) and (13a), we see that the spinons and holons behave as well defined objects with respective renormalized velocities  $v_c \approx 3\pi\zeta(3)J\delta_0/8\log^2 2$  and  $v_s \approx \pi J(1 - \delta_0)/2$ , to  $O(\delta^2)$ . Thus, the holons and spinons scatter off the impurity one by one. This will be important when we deal with low-temperature transport properties [15]. In zero field, the hole-charge is depleted in the region surrounding the defect, i.e., the impurity is screened by an amount  $\delta_i = -G(\theta)\delta_0$ , where  $G(\theta) = 1 - \pi R(\theta)/\log 2 > 0$ , while the magnetization vanishes locally, i.e., the impurity is magnetically inert. This redistribution of charge around the defect explains why the spinon- and holon-impurity phase shifts can simply be expressed in terms of the holon-holon and spinon-holon phase shifts [15].

The zero-field ratios of impurity to host susceptibilities can be shown to be  $\chi_\alpha^i/\chi_\alpha^h = [\rho_\alpha^i(q_\alpha^0) + \rho_\alpha^i(-q_\alpha^0)]/2\rho_\alpha^h(q_\alpha^0)$ , i.e., equal to the ratio of the average (over the left and right Fermi-points) density of states of the impurity to the host chain. Near half-filling, we find  $\chi_s^i/\chi_s^h \approx -\delta_0 G(\theta)$ , i.e., the local magnetic response in zero field is controlled by the screening process, while for the compressibility ratio we have  $\chi_c^i/\chi_c^h \approx G(\theta) + \delta_0^2(\pi/2 \log 2)^2[(\pi/2 \log 2)R''(\theta) + (3\pi\zeta(3)/2 \log^2 2)R(\theta) + 2R'(\theta)]$ . At finite magnetic field, the defect becomes magnetically active. For weak fields,  $h \ll J$ , and near half-filling, the Wiener-Hopf method leads to a local magnetization that is linear in  $h$ , and  $M_i/M_h \approx e^{\pi\theta} - 1 - \delta_0[e^{\pi\theta} - (\pi R(\theta)/\log 2)]$ , for  $\theta \ll |\log([e/2\pi]h)|/\pi$ . Notice that the impurity-bulk coupling is infrared finite in contrast to the Kondo problem (compare with ref. [23]). At zero temperature, we can derive simple relations between the  $h = 0$  susceptibility ratios  $\chi_\alpha^i/\chi_\alpha^h$  and their finite field counterparts,  $\chi_\alpha^i(h)/\chi_\alpha^h(h)$ , i.e.,

$$\frac{\chi_c^i(h)}{\chi_c^h(h)} = \left[1 + \frac{v_c}{v_s} \frac{Z_{cs}^2}{Z_{cc}^2}\right]^{-1} \left[\frac{\chi_c^i}{\chi_c^h} + \frac{\chi_s^i}{\chi_s^h} \frac{v_c}{v_s} \frac{Z_{cs}^2}{Z_{cc}^2}\right], \quad (14a)$$

$$\frac{\chi_s^i(h)}{\chi_s^h(h)} = \left[1 + \frac{v_c}{v_s} \left(\frac{2Z_{ss} - Z_{cs}}{2Z_{sc} - Z_{cc}}\right)^2\right]^{-1} \left[\frac{\chi_c^i}{\chi_c^h} + \frac{\chi_s^i}{\chi_s^h} \frac{v_c}{v_s} \left(\frac{2Z_{ss} - Z_{cs}}{2Z_{sc} - Z_{cc}}\right)^2\right], \quad (14b)$$

where  $v_\alpha$ ,  $\alpha = c, s$ , denotes the velocity of the  $\alpha$  pseudoparticle at the  $T = 0$  Fermi points  $\pm q_\alpha^0$  for finite  $h$  and  $Z_{\alpha\beta}$  the “dressed charge matrix” (essentially the S-matrix of the s- and c- pseudoparticles at the Fermi surface, see ref. [18]). The formulas (14a) and (14b) are universal in the sense that only the coefficients in front of the matrix elements  $Z_{\alpha\beta}$  depend on the model under consideration [15].

We have evaluated the low-temperature corrections to the thermodynamic potential to  $O(T^4)$ , from which we have inferred the low-T magnetization, entropy, susceptibilities, a.s.o. [15]. We quote the specific heat  $C_v = C_v^{(1)} + C_v^{(2)} + O(T^4)$ , with

$$\begin{aligned} C_v^{(1)} &= \frac{\pi k_B^2}{3} T \sum_\alpha \frac{1}{v_\alpha} \left(1 + \frac{1}{L} \frac{\rho_{\alpha i}(q_\alpha^0) + \rho_{\alpha i}(-q_\alpha^0)}{2\rho_{\alpha h}(q_\alpha^0)}\right), \\ C_v^{(2)} &= \frac{\pi^3 k_B^4}{3} T^3 \sum_\alpha \frac{1}{v_\alpha} \left[A_\alpha \left(1 + \frac{1}{L} \frac{\rho_{\alpha i}(q_\alpha^0) + \rho_{\alpha i}(-q_\alpha^0)}{2\rho_{\alpha h}(q_\alpha^0)}\right) \right. \\ &\quad \left. + B_\alpha \frac{\rho'_{\alpha h}(q_\alpha^0)}{\rho_{\alpha h}(q_\alpha^0)} \left(1 + \frac{1}{L} \frac{\rho'_{\alpha i}(q_\alpha^0) + \rho'_{\alpha i}(-q_\alpha^0)}{2\rho'_{\alpha h}(q_\alpha^0)}\right)\right] \end{aligned} \quad (15a)$$

$$+ C_\alpha \frac{\rho''_{\alpha h}(q_\alpha^0)}{\rho_{\alpha h}(q_\alpha^0)} \left( 1 + \frac{1}{L} \frac{\rho''_{\alpha i}(q_\alpha^0) + \rho''_{\alpha i}(-q_\alpha^0)}{2\rho''_{\alpha h}(q_\alpha^0)} \right) ] . \quad (15b)$$

The ratios of zeroth, first and second derivatives of the Fermi surface density of states of the impurity to the host chain occur in (15a) and (15b). The coefficients  $A_\alpha, B_\alpha$  and  $C_\alpha$  depend in a universal way on the Fermi surface velocity, the curvature of the spectrum and its rate of change as well as on the Scattering matrix and its higher derivatives. For the sake of brevity, we omit here these lengthy and complicated expressions [15]. The linear term in  $T$ , (15a), coincides with that inferred from the zero temperature finite size spectrum calculations [15], and can alternatively be derived from boundary conformal field theory [2]. The next order term (15b) cannot be calculated within the framework of conformal field theory. Very close to half-filling, equations (15a) and (15b) do not hold, however. At low  $T$ , when the charge carrier density  $\delta_0$  is small, a crossover takes place to a regime where the holon-liquid behaves as a non-degenerate 1D Fermi system coupled to an impurity. A crude estimate for the onset of the crossover is obtained when the width of the holon “Fermi-sea” is of order  $k_B T$ , i.e.,  $\delta_c \approx [k_B T / J]^{1/2} (4\pi^{-1} \log 2 (3\zeta(3))^{-1/2})$ . For instance, at  $T = 10\text{K}$  with an exchange of order 1 eV,  $\delta_c \approx 10^{-2}$  while at room temperature we have  $\delta_c \approx 10^{-1}$  [15]. A similar crossover takes place for the spinon liquid near the transition to the ferromagnetic state. Following ref. [9,13] we can calculate the conductance through the impurity at low- $T$ . We shall report on this elsewhere [15].

In summary, we have proposed a novel strategy to construct solvable models with impurities. The hamiltonians generated by our method contain terms that are difficult to justify on physical grounds. Yet some of those terms are, in the models considered so far, and should generically be, irrelevant to the low-energy properties of the model. We have illustrated the idea by performing explicit calculations in a simple model with a single impurity. We have derived various results that are beyond boundary conformal field theory by solving the thermodynamic Bethe ansatz equations in the high-and low-temperature limits. The transport properties of these class of models will be explored in future work.

It is a pleasure to thank P. Nozières and A. Gogolin for discussions and encouragements.

We wish to thank the organizers of a workshop on exact solutions at Aspen for the inspiring atmosphere that stimulated this work. We also gratefully acknowledge the Aspen Center for Physics for the kind hospitality during a stay in July 1994.

## REFERENCES

- [1] J. L. Cardy, Nucl. Phys. **B240**, 514 (1984); *ibid.* **B324**, 581 (1984); D. Lewellen and J. L. Cardy, Phys. Lett. **B259** 274 (1991).
- [2] I. Affleck, Nucl. Phys. **B336**, 517 (1990); I. Affleck and A. W. W. Ludwig, Phys. Rev. **B48**, 7297 (1993); see, e.g., I. Affleck in *Springer Series in Solid-State Sciences* **118**, ed. by A. Okiji and N. Kawakami (Springer-Verlag 1994) and references therein.
- [3] C. L. Kane and M. P. A. Fisher, Phys. Rev. **B46**, 15233 (1992).
- [4] V. J. Emery and S. Kivelson, Phys. Rev. **B46**, 10812 (1992).
- [5] M. Fabrizio and A. Gogolin, Phys. Rev. **B50** (1994), in press.
- [6] M. Fabrizio, A. Gogolin and P. Nozières, unpublished.
- [7] A. M. Sengupta and A. Georges, Phys. Rev. **B49**, 10020 (1994)
- [8] S. Ghoshal and A. B. Zamolodchikov, Int. J. Mod. Phys. **A9**, 3841 (1994).
- [9] P. Fendley, A. W. W. Ludwig and H. Saleur, cond-mat/9408068.
- [10] N. Andrei, H. Johannesson, Pys. Lett. **100 A**, 108 (1984).
- [11] K.-J.-B. Lee and P. Schlottmann, Phys. Rev. **B37**, 379 (1987); P. Schlottmann, J. Phys.: Condens. Matter **3** 6617 (1991).
- [12] E. Sorensen, S. Eggert and I. Affleck, J. Phys. A: Math. Gen. **26** 6757 (1993).
- [13] A. M. Tsvelick, preprint 1994.
- [14] see, e.g., L. A. Takhtajan in *Lecture Notes in Physics* **242**, ed. by B. S. Shastry, S. S. Jha, and V. Singh, (Springer, N. Y., 1985); V. E. Korepin, N. M. Bogoliubov and A. G. Izergin, *Quantum Inverse Scattering method and Correlation Functions*, Cambridge Monographs on Mathematical Physics (Cambridge University Press 1993).

- [15] P. -A. Bares, to be published.
- [16] M. Ogata, M. Luchini, S. Sorella and F. Assaad, Phys. Rev. Lett. **66**, 2388 (1991).
- [17] P. Schlottmann, Phys. Rev. **B36**, 5177 (1987).
- [18] P. A. Bares, G. Blatter and M. Ogata, Phys. Rev. **B44**, 130 (1991); P. A. Bares, J. M. P. Carmelo, J. Ferrer and P. Horsch, Phys. Rev. **B46** 14624 (1992).
- [19] N. Kawakami and S. K. Yang, J. Phys. Condens. Matter **3**, 5983 (1991).
- [20] F. Essler and V. Korepin, Phys. Rev. **B46**, 9147 (1992).
- [21] H. A. Bethe, Z. Phys. **71**, 205 (1931).
- [22] C. N. Yang and C. P. Yang, Phys. Rev. **150**, 321 (1966); *ibid.***150**, 327 (1966);*ibid.* **151**, 256 (1966).
- [23] A. M. Tsvelick and P. B. Wiegmann, Adv. Phys. **32**, 453 (1983).
- [24] M. Takahashi, Progr. Theor. Phys. **46**, 401 (1971); *ibid.* **48**, 2187 (1972);*ibid.* **50**, 1519 (1973).
- [25] To simplify the discussion, we choose a finite magnetic field. At  $h = 0$ , we have an additional branch of excitations parametrized by complex roots (two-strings) that is degenerate with  $\epsilon_s(x)$ . A complete and careful discussion of this point can be found in ref. [18].